Weed Research, 1995, Volume 35, 523–527

AUTHOR INDEX 1995

ALLEN R.L. *see* STREET *et al.*, 75
ANDERSON-TAYLOR G. *see* STREIBIG *et al.*, 215
ANDERSSON L. Effects of dose and application timing on the seed production of three weed species treated with MCPA or tribenuron-methyl, 67
ARNAUD M.C. *see* BAGONNEAUD-BERTHOME *et al.*, 25
ASCARD J. Effects of flame weeding on weed species at different developmental stages, 397

BABUT M. *see* PERRIN-GANIER *et al.*, 257
BAGONNEAUD-BERTHOME V. ARNAUD M.C. & FER A. A new experimental approach to the chemical control of *Striga* using simplified models *in vitro*, 25
BARGUTTI A. *see* GOLDWASSER *et al.*, 445
BASKIN C.C. *see* BASKIN & BASKIN, 353
BASKIN J.M. & BASKIN C.C. Variation in the annual dormancy cycle in buried seeds of the weedy winter annual *Viola arvensis*, 353
BENVENUTI S. & MACCHIA M. Effect of hypoxia on buried weed seed germination, 343
BINGHAM I.J. A comparison of the dynamics of root growth and biomass partitioning in wild oat (*Avena fatua* L.) and spring wheat, 57
BLACKSHAW R.E. & ENTZ T. Day and night temperature effects on the vegetative growth of *Erodium cicutarium*, 471
BLAIR A.M. *see* STREIBIG *et al.*, 215
BOATMAN N.D. *see* REW *et al.*, 363
BOUTSALIS P. & POWLES S.B. Resistance of dicot weeds to acetolactate synthase (ALS)-inhibiting herbicides in Australia, 149
BOYKIN D.L. *see* STREET *et al.*, 75
BRAIN P. *see* WILSON *et al.*, 265

CALLIHAN R.H. *see* MILLER & CALLIHAN, 453
CARRANZA P., SAAVEDRA M. & GARCIA-TORRES L. Competition between *Ridolfia segetum* and sunflower, 369
CASELEY J.C. *see* FISHLER *et al.*, 279
CHIKOYE D. & SWANTON C.J. Evaluation of three empirical models depicting *Ambrosia artemisiifolia* competition in white bean, 421
CHRISTENSEN S. Weed suppression ability of spring barley varieties, 241
CHUECA M.C. *see* SIXTO *et al.*, 7
CIOTOLA M., WATSON A.K. & HALLETT S.G. Discovery of an isolate of *Fusarium oxysporum* with potential to control *Striga hermonthica* in Africa, 303
CLEMENTS M. *see* WILSON *et al.*, 265
CLOUTIER D.C. *see* LEBLANC *et al.*, 511
COOMBES N. *see* LEMERLE *et al.*, 503
COTTERMAN J.C. *see* KUDSK *et al.*, 19
COX T.I. *see* POPAY *et al.*, 429

DERKSEN D.A., THOMAS A.G., LAFOND G.P., LOEPPKY H.A. & SWANTON C.J. Impact of post-emergence herbicides on weed community diversity within conservation-tillage systems, 311
DIEKKRÜGER B. *see* LÜCKE *et al.*, 333
DOLL H., HOLM U. & SOGAARD B. Effect of crop density on competition by wheat and barley with *Agrostemma githago* and other weeds, 391
DUKE S.O. *see* MADSEN *et al.*, 81

EAGLE D.J. *see* STREIBIG *et al.*, 215
EL KHYARI D. *see* LOUDYI *et al.*, 121, 225

© 1995 European Weed Research Society

ELEFTHEROHORINOS I.G. *see* HAITAS *et al.*, 1
ELEFTHEROHORINOS I.G. & KOTOULA-SYKA E. Influence of herbicide application rate and timings for post-emergence control of *Sorghum halepense* (L.) Pers. in maize, 99
ELEMO K. *see* WEBER *et al.*, 167
ENTZ T. *see* BLACKSHAW & ENTZ, 471
ESCORIAL C. *see* SIXTO *et al.*, 7

FENNER M. The effects of pre-germination chilling on subsequent growth and flowering in three arable weeds, 489
FER A. *see* BAGONNEAUD-BERTHOME *et al.*, 25
FERNANDEZ QUINTANILLA C. *see* SANCHEZ DEL ARCO *et al.*, 477
FISHLER A., CASELEY J.C., VARSANO R., NEGBI M. & RUBIN B. Factors affecting benfuresate activity against purple nutsedge (*Cyperus rotundus* L.), 279
FRIEDLÄNDER H. *see* STREIBIG *et al.*, 215
FROUD-WILLIAMS R.J. *see* Rew *et al.*, 363

GARCIA-BAUDIN J.M. *see* SIXTO *et al.*, 7
GARCIA-TORRES L. *see* CARRANZA *et al.*, 369
GHERSA C.M. & HOLT J.S. Using phenology prediction in weed management: a review, 461
GODRON M. *see* LOUDYI *et al.*, 121, 225
GOLAN S. *see* GOLDWASSER *et al.*, 445
GOLDWASSER Y., KLEIFELD Y, GOLAN S., BARGUTTI A. & RUBIN B. Dissipation of metam-sodium from the soil and its effect on the control of *Orobanche aegyptiaca*, 445
GONZALEZ PONCE R., ZANCADA C., VERDUGO M. & SALAS L. The influence of the nematode *Meloidogyne incognita* on competition between *Solanum nigrum* and tomato, 437
GUILLERM J.L. *see* LE BOURGEOIS & GUILLERM, 89
GÜNTHER P. *see* LÜCKE *et al.*, 333

HACK C. M. *see* MARSHALL *et al.*, 51
HACKER E. *see* STREIBIG *et al.*, 215
HAGGAR R.J. *see* PINO *et al.*, 141
HAITAS V.C., KOTOULA-SYKA E. & ELEFTHEROHORINOS I.G. Influence of propaquizafop application rate and time on *Sorghum halepense* (L.) Pers. control and cotton (*Gossypium hirsutum*) yield, 1
HALL J.C. *see* TAL *et al.*, 133
HALLETT S.G. *see* CIOTOLA *et al.*, 303
HEITHOLT J.J. *see* MADSEN *et al.*, 81
HILL T.A. *see* QASEM & HILL, 187
HOLLAND P.T. *see* JAMES *et al.*, 113
HOLM U. *see* DOLL *et al.*, 391
HOLT J.S. *see* GHERSA & HOLT, 461
HSAIO A.I. *see* SHUMA *et al.*, 249

INGLE A. *see* POPAY *et al.*, 429
IRIONDO J.M. *see* PEREZ-GARCIA *et al.*, 495
IWANZIK W. *see* STREIBIG *et al.*, 215

JAMES T.K., KLAFFENBACH P., HOLLAND P.T. & RAHMAN A. Degradation of primisulfuron-methyl and metasulfuron-methyl in soil, 113
JENSEN J.E. *see* MADSEN & JENSEN, 105
JOHNSON G.A., MORTENSEN D.A. & MARTIN A.R. A simulation of herbicide use based on weed spatial distribution, 197

KEMPENAAR C. & SCHNIEDERS B.J. A method to obtain fast and uniform emergence of weeds for field experiments, 385
KERR R. *see* POPAY *et al.*, 429

KIRKWOOD R.C. *see* MARSHALL *et al.*, 51
KLAFFENBACH P. *see* JAMES *et al.*, 113
KLEIFELD Y. *see* GOLDWASSER *et al.*, 445
KNEŽEVIĆ S.Ž., WEISE S.F., & SWANTON C.J. Comparison of empirical models depicting density of *Amaranthus retroflexus* L. and relative leaf area as predictors of yield loss in maize (*Zea mays* L.), 207
KORPELAINEN H. Geographical differentiation in allozyme variation in Rumex acetosella subspecies acetosella and angiocarpus, 413
KOTOULA-SYKA E. *see* ELEFTHEROHORINOS & KOTOULA-SYKA, 99
KOTOULA-SYKA E. *see* HAITAS *et al.*, 1
KUDSK P. *see* STREIBIG *et al.*, 215
KUDSK P., MATHIASSEN S.K. & COTTERMAN J.C. Sulfonylurea resistance in *Stellaria media* (L.) Vill, 19

LABHART C. *see* STREIBIG *et al.*, 215
LAFOND G.P. *see* DERKSEN *et al.*, 311
LAGOKE S.T.O. *see* WEBER *et al.*, 167
LE BOURGEOIS T. & GUILLERM J.L. Etendue de distribution et degré d'infestation des adventices dans la rotation cotonnière au Nord-Cameroun, 89
LEBLANC M.L., CLOUTIER D.C. & LEROUX G.D. Réduction de l'utilisation des herbicides dans le maïs-grain par une application d'herbicides en bandes combinée à des sarclages mécaniques, 511
LEMERLE D., VERBEEK B. & COOMBES N. Losses in grain yield of winter crops from *Lolium rigidum* competition depend on crop species, cultivar and season, 503
LEOPPKY H.A. *see* DERKSEN *et al.*, 311
LEROUX G.D. *see* LEBLANC *et al.*, 511
LEROUX G.D. *see* MEKKI & LEROUX, 377
LOUDYI M.C., GODRON M. & EL KHYARI D. Influence des variables écologiques sur la distribution des mauvaises herbes des cultures du Saïs (Maroc central), 225
LOUDYI M.C., GODRON M. & EL KHYARI D. Profils écologiques et liaisons interspécifiques. Deux approches complémentaires appliquées à l'étude des groupements de mauvaises herbes des cultures du Saïs (Maroc central), 121
LÜCKE A., GÜNTHER P., DIEKKRÜGER B, PESTEMER W. & RICHTER O. Implementation of a herbicide simulation model in a geographical information system with an example of a site-specific application, 333
LUSCOMBE B.M. *see* STREIBIG *et al.*, 215

MACCHIA M. *see* BENVENUTI & MACCHIA, 343
MADAFIGLIO G. *see* STREIBIG *et al.*, 215
MADDEN A.D. An assessment, using a modelling approach, of inbreeding as a possible cause of reduced competitiveness in triazine-resistant weeds, 289
MADSEN K.H., HEITHOLT J.J., DUKE, S.O. SMEDA R.J. & STREIBIG J.C. Photosynthetic parameters in glyphosphate-treated sugarbeet (*Beta vulgaris* L.), 81
MADSEN K.H. & JENSEN J.E. Weed control in glyphosphate-tolerant sugarbeet (*Beta vulgaris* L.), 105
MARSHALL G., HACK C. M. & KIRKWOOD R.C. Volunteer barley interference in fibre flax (*Linum usitatissimum* L.), 51
MARTIN A.R. *see* JOHNSON *et al.*, 197
MARTINEZ-LABORDE J.B. *see* PEREZ-GARCIA *et al.*, 495
MASALLES R.M. *see* PINO *et al.*, 141
MASALLES R.M. *see* SANS & MASALLES, 321
MATHIASSEN S.K. *see* KUDSK *et al.*, 19
MEKKI M. & LEROUX G.D. Foliar absorption and translocation of nicosulfuron and rimsulfuron in five annual weed species, 377
MELANDER B. Impact of drilling date on *Apera spica-venti* L. and *Alopecurus myosuroides* Huds. in winter cereals, 157
MERSIE W. Absorption, translocation and fate of propyzamide in witloof chicory (Cichorium intybus) and common amaranth (*Amaranthus retroflexus*), 15

MILLER T.W. & CALLIHAN R.H. Interference between triazine-resistant *Brassica napus* and *Panicum miliaceum*, 453
MORTENSEN D.A. *see* JOHNSON *et al.*, 197

NEGBI M. *see* FISHLER *et al.*, 279
NEL P.C. *see* STREIBIG *et al.*, 215

PEREZ-GARCIA F., IRIONDO J.M. & MARTINEZ-LABORDE J.B. Germination behaviour in seeds of *Diplotaxis erucoides* and *D. virgata*, 495
PERRIN-GANIER C., SCHIAVON M., PORTAL J.-M. & BABUT M. Dégradation de l'isoproturon et disponibilité de ses résidus dans le sol, 257
PESTEMER W. *see* LÜCKE *et al.*, 333
PESTEMER W. *see* STREIBIG *et al.*, 215
PINO J., HAGGAR R.J., SANS F.X., MASALLES R.M. & SACKVILLE-HAMILTON R.N. Clonal growth and fragment regeneration of *Rumex obtusifolis* L., 141
POPAY A.I., COX T.I., INGLE A. & KERR R. Seasonal emergence of weeds in cultivated soil in New Zealand, 429
PORTAL J.-M. *see* PERRIN-GANIER *et al.*, 257
POWLES S.B. *see* BOUTSALIS & POWLES, 149

QASEM J.R. & HILL T.A. Growth, development and nutrient accumulation in *Senecio vulgaris* L. and *Chenopodium album* L., 187
QASEM J.R. The allelopathic effect of three *Amaranthus* spp.(pigweeds) on wheat (*Triticum durum*), 41
QUICK W.A. *see* SHUMA *et al.*, 249

RAHMAN A. *see* JAMES *et al.*, 113
RAHMAN A. *see* STREIBIG *et al.*, 215
RAJU M.V.S. *see* SHUMA *et al.*, 249
RETZLAFF G. *see* STREIBIG *et al.*, 215
REW L.J., FROUD-WILLIAMS R.J. & BOATMAN N.D. The effect of nitrogen, plant density and competition between *Bromus sterilis* and three perennial grasses: the implications for boundary strip management, 363
RICHTER O. *see* LÜCKE *et al.*, 333
ROCHA F. & WALKER A. Simulation of the persistance of atrazine in soil at different sites in Portugal, 179
ROLA J. *see* STREIBIG *et al.*, 215
RUBIN B. *see* FISHLER *et al.*, 279
RUBIN B. *see* GOLDWASSER *et al.*, 445

SAAVEDRA M. *see* CARRANZA *et al.*, 369
SABADIE J. Alcoolyse et hydrolyse chimique du chlorimuron éthyle, 33
SABADIE J. Dégradation du chlorimuron-éthyle déposé sur divers minéraux et sur acides humiques, 295
SACKVILLE-HAMILTON R.N. *see* Pino *et al.*, 141
SALAS L. *see* GONZALEZ PONCE *et al.*, 437
SANCHEZ DEL ARCO M.J., TORNER C. & FERNANDEZ QUINTANILLA C. Seed dynamics in populations of *Avena sterilis* ssp. *ludoviciana*, 477
SANS F.X. *see* PINO *et al.*, 141
SANS F.X. & MASALLES R.M. Phenological patterns in an arable land weed community related to disturbance, 321
SCHIAVON M. *see* PERRIN-GANIER *et al.*, 257
SCHNIEDERS B.J. *see* KEMPENAAR & SCHNIEDERS, 385
SHUMA J.M., QUICK W.A., RAJU M.V.S. & HSAIO A.I. Germination of seeds from plants of *Avena fatua* L. treated with glyphosate, 249

SILVELA L. *see* SIXTO *et al.*, 7
SIXTO H., SILVELA L., ESCORIAL C., GARCIA-BAUDIN J.M. & CHUECA M.C. On the inheritance of tolerance to chlorotoluron application in wheat using a very efficient score test, 7
SMEDA R.J. *see* MADSEN *et al.*, 81
SOGAARD B. *see* DOLL *et al.*, 391
STEFANOVIC L. *see* STREIBIG *et al.*, 215
STEPHENS E. *see* WILSON *et al.*, 265
STEPHENSON G.R. *see* TAL *et al.*, 133
STRAATHOF H.J.M. *see* STREIBIG *et al.*, 215
STREET J.E., TERESIAK H., BOYKIN D.L. & ALLEN R.L. Interaction between timings and doses of quinclorac in rice, 75
STREIBIG J.C. *see* MADSEN *et al.*, 81
STREIBIG J.C., WALKER A., BLAIR A.M., ANDERSON-TAYLOR G., EAGLE D.J., FREIDLÄNDER H., HACKER E., IWANZIK W., KUDSK P., LABHART C., LUSCOMBE B.M., MADAFIGLIO G., NEL P.C., PESTEMER W., RAHMANA., RETZLAFF G., ROLA J., STEFANOVIC L., STRAATHOF H.J.M. &THIES E.P. Variability of bioassays with metsulfuron-methyl in soil, 215
SWANTON C.J. *see* CHIKOYE & SWANTON, 421
SWANTON C.J. *see* DERKSEN *et al.*, 311
SWANTON C.J. *see* KNEŽEVIĆ *et al.*, 207

TAL J.A., HALL J.C. & STEPHENSON G.R. Non-enzymatic conjugation of fenoxaprop-ethyl with glutathione and cysteine in several grass species, 133
TERESIAK H. *see* STREET *et al.*, 75
THIES E.P. *see* STREIBIG *et al.*, 215
THOMAS A.G. *see* DERKSEN *et al.*, 311
TORNER C. *see* SANCHEZ DEL ARCO *et al.*, 477

VARSANO R. *see* FISHLER *et al.*, 279
VERBEEK B. *see* LEMERLE *et al.*, 503
VERDUGO M. *see* GONZALEZ PONCE *et al.*, 437

WALKER A. *see* ROCHA & WALKER, 179
WALKER A. *see* STREIBIG *et al.*, 215
WATSON A.K. *see* CIOTOLA *et al.*, 303
WEBER G., ELEMO K. & LAGOKE S.T.O. Weed communities in intensified cereal-based cropping systems of the northern Guinea savanna, 167
WEISE S.F. *see* KNEŽEVIĆ *et al.*, 207
WILSON B.J., WRIGHT K.J., BRAIN P., CLEMENTS M. & STEPHENS E. Predicting the competitive effects of weed and crop density on weed biomass, weed seed production and crop yield in wheat, 265
WRIGHT K.J. *see* WILSON *et al.*, 265

ZANCADA C. *see* GONZALEZ PONCE *et al.*, 437